



## Food Supplement with Sucrosomial® Iron Bottle

BOTTLE 30ml + STICK 2,5g 

**Indications:** Sideral® Gocce Forte is a food supplement with Sucrosomial® Iron (Sideral® r.m.), useful to cover dietary deficiencies or an increased demand for said nutrient. Sucrosomial® Iron can pass the gastric environment intact to be directly absorbed in the intestine, thus avoiding eventual gastric disorders.

Iron is useful for the correct cognitive development of children. Iron contributes to normal formation of red blood cells and haemoglobin, to normal oxygen transport in the body, to normal cognitive function, to normal energy metabolism, to the normal function of the immune system and to the reduction of tiredness and fatigue. Iron has a role in the process of cell division.

**Stick Ingredients:** Sideral® r.m. - Sucrosomial® Iron (iron pyrophosphate, sucrose esters of fatty acids, sunflower lecithin spray-dried on rice flour and tricalcium phosphate, pregelatinized rice starch), maltodextrins. Gluten free.

**Bottle Ingredients:** Demineralized water, sucrose, maltodextrins, preservatives: potassium sorbate, sodium benzoate; flavour; thickening agent: xanthan gum; acidifying agent: citric acid; emulsifier: sucrose esters of fatty acids. Gluten free.

**Warnings:** Do not take more than the advised daily dosage. Keep out of the reach of children under 3 years of age. Food supplements should not be considered as a replacement for a varied, balanced diet and a healthy lifestyle. Use only after reconstitution.

**How to reconstitute:** 1) Open the bottle. Be careful not to waste the cap. 2) Put the powder inside the bottle. 3) Put the cap on the bottle (do not use the dropper) and close it well, then let it sit for one minute. 4) Shake the bottle vigorously. 5) Shake well before each use.

Mean Contents	Per daily dose (1ml)	%NRVs
Iron	14 mg	100%

%NRVs: Nutrient Reference Values Reg. 1169/2011

**Storage:** Store the product at room temperature (do not exceed 25°C) avoiding exposure to localized heat sources, sunlight and water. After reconstitution store at 4°C. Use within one month of reconstitution.

The presence of agglomerates immediately after reconstitution is due to the presence of anticaking agents inside the powder. It tends to disappear within 10 minutes after the reconstitution. Therefore, we suggest to reconstitute the product a few minutes before use.

**Directions:** Advised dosage is 1ml a day, equal to about 12 drops, or as suggested by the doctor. Use the dropper included in the package. The product can be taken directly in the mouth or dissolved in a glass of water at room temperature, at any time of the day. Shake well before each use after reconstitution.

**The presence of agglomerates immediately after reconstitution is due to the presence of anticaking agents inside the powder. It tends to disappear within 10 minutes after the reconstitution. Therefore, we suggest to reconstitute the product a few minutes before use.**

Sucrosomial® Iron is a registered trademark.  
Sucrosome® Iron is a registered trademark.  
Sideral® is a registered trademark.  
Sideral® r.m. is a patented ingredient based on Sucrosomial® Iron.

 JUNIAPHARMA

Produced and packaged in the facility based in Via Ariosto 50/60 - 20090 Trezzano S/N (Milano) - Italy  
on behalf of Junia Pharma S.r.l. Via Campodavola, 1 - 56122 Pisa (PI) - Italy  
[www.juniapharma.it](http://www.juniapharma.it)

